Registration form

Tick Control CEU Training 48 HOUR RUSH ORDER PROCESSING FEE ADDITIONAL \$50.00 Rush service does not include overnight delivery or FedEx fees.

Name	Signature	
Address:		
City	State	_Zip_
Phone: Home ()	Work ()	
Fax ()	Email	
License #	Ехр. І	Date_
Class/Grade		
Please circle/check which c	ertification you are applying the course C	EU's
Commercial Applicator Re	esidential Applicator Industrial Applicator	
Pesticide Handler Agricultu	ural Applicator Adviser Other	
F	Technical Learning College PO Box 3060, Chino Valley, AZ 86323-3060 Fax (928) 272-0747 e-mail info@tlch2o.c (928) 468-0665 Toll Free (866) 557-1746	om

We will stop mailing the certificate of completion we need your e-mail address. We will e-mail the certificate to you.

DISCLAIMER NOTICE

I understand that it is my responsibility to ensure that this CEU course is either approved or accepted in my State for CEU credit. I understand State laws and rules change on a frequent basis and I believe this course is currently accepted in my State for CEU or contact hour credit, if it is not, I will not hold Technical Learning College responsible. I fully understand that this type of study program deals with dangerous, changing conditions and various laws and that I will not hold Technical Learning College, Technical Learning Consultants, Inc. (TLC) liable in any fashion for any errors, omissions, advice, suggestions or neglect contained in this CEU education training course or for any violation or injury, death, neglect, damage or loss of your license or certification caused in any fashion by this CEU education training or course material suggestion or error or my lack of submitting paperwork. It is my responsibility to call or contact TLC if I need help or assistance and double-check to ensure my registration page and assignment has been received and graded. It is my responsibility to ensure all information is correct and to abide with all rules and regulations.

State Approval Listing Link, check to see if your State accepts or has pre-approved this course. Not all States are listed. Not all courses are listed.

State Approval Listing URL...

http://www.ABCTLC.com/downloads/PDF/CEU%20State%20Approvals.pdf

You can obtain a printed version of the course manual from TLC for an additional \$169.95 plus shipping charges.

AFFIDAVIT OF EXAM COMPLETION

I affirm that I personally completed the entire text of the course. I also affirm that I completed the exam without assistance from any outside source. I understand that it is my responsibility to file or maintain my certificate of completion as required by the state or by the designation organization.

Grading Information

In order to maintain the integrity of our courses we do not distribute test scores, percentages or questions missed. Our exams are based upon pass/fail criteria with the benchmark for successful completion set at 70%. Once you pass the exam, your record will reflect a successful completion and a certificate will be issued to you.

For security purposes, please fax or e-mail a copy of your driver's license and always call us to confirm we've received your assignment and to confirm your identity.

Thank you...

CUSTOMER SERVICE RESPONSE CARD

Tick Control Training Course

	PLETE THIS FO E ANSWER IN T		ING THE NUMBER LOW.	OF THE
Please Very Easy	rate the difficulty 0 1	of your course 2 3 4	e. 5 Very Diffic	ult
Please Very Easy	rate the difficulty 0 1	of the testing 2 3 4	process. 5 Very Diffic	ult
			exam to your actual fi 5 Very Differ	
How did	d you hear about	this Course?		
What w	ould you do to in	nprove the Cou	ırse?	
Poor	the price of the Fair Ave our customer se	rage Goo	odGreat	

CERTIFICATION OF COURSE PROCTOR

Technical Learning College requires that our students who takes a correspondence or home study program course must pass a proctored course reading, quiz and final examination. The proctor must complete and provide to the school a certification form approved by the commission for each examination administered by the proctor.

Instructions. When a student completes the course work, fill out the blanks in this section and provide the form to the proctor with the examination.

Name of Course:_____

Name of Licensee:_____

Instructions to Proctor. After an examination is administered, complete and return this certification and examination to the school in a sealed exam packet or in pdf format.

I certify that:

- 1. I am a disinterested third party in the administration of this examination. I am not related by blood, marriage or any other relationship to the licensee which would influence me from properly administering the examination.
- 2. The licensee showed me positive photo identification prior to completing the examination.
- The enclosed examination was administered under my supervision on _____. The licensee received no assistance and had no access to books, notes or reference material.
- 4. I have not permitted the examination to be compromised, copied, or recorded in any way or by any method.
- 5. Provide an estimate of the amount of time the student took to complete the assignment.

Time to complete the entire course and final exam.

Notation of any problem or concerns:

Name and Telephone of Proctor (please print):

Signature of Proctor

Tick Control Answer Key

Name_____

Phone#_____

You are solely responsible in ensuring this course is accepted by your State for credit. No refunds. Did you check with your State agency to ensure this course is accepted for credit?

Method of Course acceptance confirmation. Please fill this section

Website __ Telephone Call___ Email____ Spoke to_____

Did you receive the approval number, if applicable?

What is the course approval number, if applicable?

You are responsible to ensure that TLC receives the Assignment and Registration Key. Please call us to ensure that we received it.

Multiple Choice. Pick only one answer per question.

Circle or Mark off or Bold the answer. Please circle the number of the assignment version 1 or 2 or 3 or 4 or 5

Complete all the Topical Sections before submitting the answer key

Topic 1 - Introduction to Ticks

1. АВСД	5. A B C D	9. A B C D
2. A B C D	6. A B C D	10. A B C D
3. A B C D	7. A B C D	
4. A B C D	8. A B C D	
Topic 2 - Tick Identification	Section	
1. A B C D	5. A B C D	9. A B C D
2. A B C D	6. A B C D	10. A B C D
3. A B C D	7. A B C D	
4. A B C D	8. A B C D	

Topic 3 - Dangers of Ticks		
1. A B C D	5. A B C D	9. A B C D
2. A B C D	6. A B C D	10. a b c d
3. A B C D	7. A B C D	
4. A B C D	8. A B C D	
Topic 4 – Related Blood-feeding	Insects	
Topic 4 – Related Blood-feeding 1. A B C D	Insects 5. A B C D	9. A B C D
		9. АВСД 10. АВСД
1. A B C D	5. A B C D	
1. АВСД 2. АВСД	5. A B C D 6. A B C D	

I understand that I am 100 percent responsible to ensure that TLC receives the Assignment and Registration Key. I understand that TLC has a zero tolerance towards not following their rules, cheating or hostility towards staff or instructors. I need to complete the entire assignment for credit. My exam was proctored. There is no credit for partial assignment completion. I will contact TLC if I do not hear back from them within 2 days of assignment submission. I will not hold TLC liable for any errors, injury, death or non-compliance with rules. I will abide with all federal and state rules and rules found on page 2 and 7. I will forfeit my purchase costs and will not receive credit or a refund if I do not abide with TLC's rules.

California DPR Requirement

The Assignment must be submitted to TLC by December 27 in order to be submitted to DPR by the 31st. If it is late, you will be penalized \$50 per day.

Please sign that you understand and will abide with TLC's Rules and disclaimer page 2 and 7

Signature

Email info@tlch2o.com

Important Information about this Course (Disclaimer Notice)

This CEU course has been prepared to educate pesticide applicators and operators in general safety awareness of dealing with the often-complex and various pesticide treatment sprays, devices, methods, and applications. This course (manual) will cover general laws, regulations, required procedures and accepted policies relating to the use of pesticides and herbicides. It should be noted, however, that the regulation of pesticides and hazardous materials is an ongoing process and subject to change over time. For this reason, a list of resources is provided to assist in obtaining the most up-to-date information on various subjects. This manual is a not a guidance document for applicators or operators who are involved with pesticides. It is not designed to meet the requirements of the United States Environmental Protection Agency or your local State environmental protection agency or health department. This course manual will provide general pesticide safety awareness and should not be used as a basis for pesticide treatment method/device guidance. This document is not a detailed pesticide informational manual or a source or remedy for poison control.

Technical Learning College or Technical Learning Consultants, Inc. makes no warranty, guarantee or representation as to the absolute correctness or appropriateness of the information in this manual and assumes no responsibility in connection with the implementation of this information. It cannot be assumed that this manual contains all measures and concepts required for specific conditions or circumstances. This document should be used for educational purposes only and is not considered a legal document. Pesticides are poisonous. Always read and carefully follow all precautions and safety recommendations given on the container label. Store all chemicals in the original labeled containers in a locked cabinet or shed, away from food or feeds, and out of the reach of children, unauthorized persons, pets, and livestock.

Confine chemicals to the property or plants being treated. Avoid drift onto neighboring properties, especially gardens containing fruits and/or vegetables ready to be picked. Dispose of empty containers carefully. Follow label instructions for disposal. Never reuse containers. Make sure empty containers are not accessible to children or animals. Never dispose of containers where they may contaminate water supplies or natural waterways. Do not pour down sink or toilet. Consult your county agricultural commissioner for correct ways of disposing of excess pesticides. You should never burn pesticide containers.

Individuals who are responsible for pesticide storage, mixing and application should obtain and comply with the most recent federal, state, and local regulations relevant to these sites and are urged to consult with the EPA and other appropriate federal, state and local agencies.

USE PESTICIDES WISELY: ALWAYS READ THE ENTIRE PESTICIDE LABEL CAREFULLY, FOLLOW ALL MIXING AND APPLICATION INSTRUCTIONS AND WEAR ALL RECOMMENDED PERSONAL PROTECTIVE GEAR AND CLOTHING. CONTACT YOUR STATE DEPARTMENT OF AGRICULTURE FOR ANY ADDITIONAL PESTICIDE USE REQUIREMENTS, RESTRICTIONS OR RECOMMENDATIONS.

NOTICE: MENTION OF PESTICIDE PRODUCTS IN THIS COURSE DOES NOT CONSTITUTE ENDORSEMENT OF ANY MATERIAL OR HERB OR HERBAL SUPPLEMENT. ALWAYS FOLLOW THE PRODUCT'S LABEL INSTRUCTIONS.

NOTICE

I understand that it is my responsibility to ensure that this CEU course is either approved or accepted in my State for CEU credit. I understand State laws and rules change on a frequent basis and I believe this course is currently accepted in my State for CEU or contact hour credit, if it is not, I will not hold Technical Learning College responsible. I also understand that this type of study program deals with dangerous conditions and that I will not hold Technical Learning College, Technical Learning Consultants, Inc. (TLC) liable for any errors or omissions or advice contained in this CEU education training course or for any violation or injury caused by this CEU education training course material. I will call or contact TLC if I need help or assistance and double-check to ensure my registration page and assignment has been received and graded.

Grading Information

In order to maintain the integrity of our courses we do not distribute test scores, percentages or questions missed. Our exams are based upon pass/fail criteria with the benchmark for successful completion set at 70%. Once you pass the exam, your record will reflect a successful completion and a certificate will be issued to you.

INSTRUCTIONS

1. We will require all students to fax or e-mail a copy of their driver's license with the registration form.

2. You will need to pick one of the following four assignments to complete. This selection process is based upon your last name.

Assignment for Last Names *If your last name...*

A-G Assignment #1 pages 9-14

- H-M Assignment #2 Pages 15-20
- N-S Assignment #3 Pages 21-26
- T-Z Assignment #4 Pages 27-32

Alternative Assignment #5 for repeat students Pages 33-37

These exams are frequently rotated. Complete all topics before submitting the answers key.

Rush Grading Service

If you need this assignment graded and the results mailed to you within a 48-hour period, prepare to pay an additional rush service handling fee of \$50.00. This fee may not cover postage costs. If you need this service, simply write RUSH on the top of your Registration Form. We will place you in the front of the grading and processing line.

When Finished with Your Assignment

REQUIRED DOCUMENTS

Please scan the **Registration Page**, **Answer Key**, **Survey and Driver's License** and email it to <u>info@TLCH2O.com</u>.

IPhone Scanning Instructions

If you are unable to scan, take a photo of these documents with your **iPhone** and send these photos to TLC, <u>info@TLCH2O.com</u>.

FAX

If you are unable to scan and email, please fax these to TLC, if you fax, call to confirm that we received your paperwork. **(928) 468-0675**

Tick Control CEU Training Awareness Assignment #1 For Students Names A-G

You will have 90 days from the start of this course to have successfully passed this assignment with a score of 70 %. You may e mail the answers to TLC, info@tlch2o.com or fax the answers to TLC, (928) 272-0747.

This assignment is available to you in a Word Format on TLC's Website. You can find online assistance for this course on the in the Search function on Adobe Acrobat PDF to help find the answers. Student Services (928) 468-0665.

Topic 1 Introduction to Ticks

Multiple choice. Please select one answer only per question. No trick questions.

(S) means the answer may be plural or singular in nature. Or means either answer may work.

1. More than 800 species of ticks inhabit the planet. They are second only to mosquitoes as vectors of human disease,

- A. Including parasitic mechanisms C. Both infectious and toxic
- B. Causing allergic reaction(s) D. None of the above

2. Ixodidae or Hard Ticks >700 species are distinguished from the Argasidae by the presence of a or hard shield.

- C. Scutum A. Idiosoma
- B. Capitulum (head) D. None of the above

Life cycle and reproduction

ticks undergo three primary stages of development: larval, nymphal, 3.

and adult.

- A. Only Argasidae or Argasid C. Both ixodid and argasid
- B. Only Dermacentor
- D. None of the above

Ixodidae

- 4. Ixodid ticks require three hosts, and their life cycle takes at least one year to complete. Up to 3,000 eggs are laid on the ground by an adult female tick.
- A. 100 C. 500
- B. 3,000 D. None of the above

5. All ticks have an incomplete metamorphosis: after hatching from the egg a series of similar stages (instars) develop from a , to eight legged nymph and then a sexually developed eight legged adult.

- A. Six legged larva C. Eight legged larva
- B. Seven instar D. None of the above

6. Between each stage there is a molt (ecdysis) which enables the developing tick to expand within a new

- C. External skeleton A. Idiosoma
- B. Haller's organ D. None of the above

7. The family ______ contains the important genera Amblyomma, Dermacentor, Haemaphysalis, Hyalomma, Ixodes, Margaropus, and Rhipicephalus. Also the important boophilid ticks, formerly of the genus Boophilus, are now classified as a sub-genus within the genus Rhipicephalus.

A. Ornithodoros C. Dermacentor

B. Ixodidae D. None of the above

- 8. The cement serves to hold the in place while the tick feeds.
- A. IdiosomaC. MouthpartsB. CapitulumD. None of the above

on larval and nymphal ticks are small with less penetration and 9. produce a smaller host reaction.

- A. Idiosoma C. Mouthparts
- B. Hypostome D. None of the above

10. Adult Ixodes and ______ticks have long mouthparts that can reach the sub dermal layer of skin, produce a larger reaction, and make the tick harder to remove.

- A. Argasidae or Argasid C. Dermacentor
- B. Amblyomma D. None of the above

Please complete the entire assignment before submitting the answer key

Topic 2 Tick Identification Section

Deer Tick Life Cycle

- 1. The deer tick passes through four life stages (egg, larva, nymph, adult), over a
- A. Two month periodC. Two year periodB. Three month periodD. None of the above

Egg to Larvae

2. Eggs are fertilized in the fall and deposited in leaf litter the following .

- A. Summer C. Spring
- B. Month D. None of the above

3. The larvae then drop off their host into the leaf litter where they molt into the next stage,

- the nymph, remaining dormant until the following ______.
- A. SummerC. SpringB. MonthD. None of the above

Larvae to Nymph

4. During the spring and early summer of the next year the nymphs end their dormancy and begin to seek a host. ______ are commonly found on the forest floor in leaf litter and on low lying vegetation.

- A. Nymph(s)
- C. Females
- B. Seven instars D. None of the above

Nymph to Adult

5. Over the next few months the nymph molts into the larger adult tick, which emerges in fall, with a peak in October through November. find and feed on a host, then

the females lay eggs sometime after feeding.

A. Both male and female adults C. Larvae

B. Seven instars D. None of the above

Adult Ticks

- 6. In the fall of the second year, nymphs molt into adult ticks. Female adults are and larger than males.
- A. Red or orange C. Black
- B. Black and red D. None of the above

7. As female ticks feed over the course of several days, their bodies slowly enlarge with blood (engorge). Adult females infected with disease agents as may transmit disease during this feeding.

- A. Both male and female adults C. Several nymphal stages
- B. Larvae or nymphs
- D. None of the above

8. ticks attach, but do not feed or become engorged. Because the adult males do not take a blood meal, they do not transmit Lyme disease, human anaplasmosis, or babesiosis.

- A. Nymph(s) C. The adult female
- D. None of the above B. Male

Lone Star Tick Amblyomma americanum eggs, which are deposited under leaf and

9. Each female produces

soil litter in middle to late spring. C. 3.000-8.000 A. 300-800

B. 30,000-80,000 D. None of the above

Winter Tick Dermacentor albipictus

10. is found throughout North America. It is widely distributed throughout California, but populations are concentrated around the central coastal and sierra foothill areas. It primarily feeds on horses and deer from fall through early spring. Heavy infestations of horses may cause emaciation and anemia (Furman and Loomis 1984).

A. This two host tick C. This one host tick

B. This no host tick D. None of the above

Topic 3 Dangers of Ticks

1. Ticks may cause in humans that is reversible when the ticks are removed. Symptoms include paralysis of the arms and legs, followed by a general paralysis, which can be fatal if not reversed.

- A. Allergic Reaction C. Local infection
- B. Paralysis D. None of the above

2. The victim may recover completely within a of the removal of the tick.

- A. Few weeks C. Few hours
- B. 1 week D. None of the above

3. The paralysis may be cau humans when a tick feeds.	ised by a	transmitted to
A. Blood C. Ge	rm	
B. Salivary toxin D. No	ne of the above	
 The highest incidence of the Columbia, Canada, and the the A. United States B. Mexico 	tick paralysis in North America occ northwestern C. Washington D. None of the above	curs near the border of British
6. The two most important ti and Rocky Mountain spottedA. Lyme diseaseB. BabesiosisC. LocD. No	cal infections	tates are
rash, which may develop a c	se is usually characterized by the c haracteristic clear central area (" bite, often in the area around the cal infection ne of the above	"),
	ymptom of Rocky Mountain spotte C. One to two years D. None of the above	
the development of tick para A. Disease symptoms	nt the transmission of lysis is the prompt removal of ticks C. Hepatitis D. None of the above	and
	hment; some back-and-forth wiggl C. Retract or pull	
Topic 4 Related Bloo Grain mite 1. Grain mites are reported to under various names as " and "copra itch." A. Grocers' itch	d-feeding Insects to have been the cause of mild der ," "vanillism" (from i C. Rickettsial pox	rmatitis in humans, known nfestations on vanilla beans),

B. Rat-mite dermatitis D. None of the above

Bed Buas

2. Cimicidae or bed bugs (sometimes bedbugs), are small parasitic insects. The most common type is Cimex lectularius.

A. True B. False

Life Stages

3. Newly hatched nymphs are translucent, lighter in color and become browner as they molt and reach maturity. Bed bugs may be mistaken for other insects such as

- ____, or vice-versa. A. Fleas or ticks
 - C. Booklice and carpet beetles
- B. Rodents or Rats D. None of the above

Pvrethroids

4. To mimic the insecticidal activity of the natural compound pyrethrum another class of pesticides, pyrethroid pesticides, has been developed. These are , which is a sodium channel modulators, and are much less acutely toxic than organophosphates and carbamates. Compounds in this group are often applied against household pests.

A. Beneficial C. Non-systemic insecticide

D. None of the above B. Non-persistent

Pyrethroids include:

Bifenthrin, Beta-Cyfluthrin, Cyfluthrin, Cypermethrin, 5. Cyphenothrin, Deltamethrin, Esfenvalerate, Fenpropathrin, Tau-Fluvalinate, Lambda-Cyhalothrin, Gamma Cyhalothrin, Imiprothrin, 1RS cis-Permethrin, Permethrin, Prallethrin, Resmethrin, Sumithrin (d-phenothrin), Tefluthrin, Tetramethrin, Tralomethrin, and Zeta-Cypermethrin

A. Beneficial

C. Non-systemic insecticide D. None of the above

Permethrin

General Information

B. Allethrin stereoisomers

6. Permethrin is a ______. It is available in dusts, emulsifiable concentrates, smokes, ULV concentrates, and wettable-powder formulations.

- A. An insect growth regulator (IGR) C. A broad-spectrum pyrethroid insecticide
- B. Stereoisomers

D. None of the above

Common Flea Treatment Pesticides Propoxur

7. Propoxur (Baygon®) is a ______insecticide and was introduced in 1959. Propoxur is a non-systemic insecticide with a fast knockdown and long residual effect used against turf, forestry, and household pests and fleas. It is also used in pest control for other domestic animals.

A. Carbamate

- C. Isopropanol
- B. Sodium channel modulators D. None of the above

Methoprene IGR Treatment

8. Methoprene is ______ with activity against a variety of insect species including horn flies, mosquitoes, beetles, tobacco moths, sciarid flies, fleas (eggs and larvae), fire ants, pharoah ants, midge flies and Indian meal moths. Controlling some of these insects, methoprene is used in the production of a number of foods including meat, milk, mushrooms, peanuts, rice and cereals. It also has several uses on domestic animals (pets) for controlling fleas.

A. Not a pyrethrin and permethrin B. A stereoisomer

C. An insect growth regulator (IGR) D. None of the above

The three types of lice that live on humans are:

9. Only the _______ is known to spread disease. Lice infestations (pediculosis and pthiriasis) are spread most commonly by close person-to-person contact.

A. Lice C. Body louse

B. Flea D. None of the above

Lindane

10. Lindane has been associated with ______suffered both by people being treated and by people applying the treatment. It is also a troublesome pollutant of wastewater and requires special treatment to be removed. While lindane is still available by prescription, pyrethrin and permethrin are safer, more effective, and less polluting than lindane. A. A variety of adverse reactions C. An insect growth regulators

A. A variety of adverse reaction B. Long residual effects

D. None of the above

California DPR Requirement

The Assignment must be submitted to TLC by December 27 in order to be submitted to DPR by the 31st. If it is late, you will be penalized \$50 per day.

When Finished with Your Assignment

REQUIRED DOCUMENTS

Please scan the **Registration Page**, **Proctoring report**, **Answer Key**, **Survey and Driver's License** and email it to <u>info@TLCH2O.com</u>.

IPhone Scanning Instructions

If you are unable to scan, take a photo of these documents with your **iPhone** and send these photos to TLC, <u>info@TLCH2O.com</u>.

FAX

If you are unable to scan and email, please fax these to TLC, if you fax, call to confirm that we received your paperwork. **(928) 468-0675**

California DPR Requirement

The Assignment must be submitted to TLC by December 27 in order to be submitted to DPR by the 31st. If it is late, you will be penalized \$50 per day.

Tick Control CEU Training Awareness Assignment #2 For Students Names H-M

You will have 90 days from the start of this course to have successfully passed this assignment with a score of 70 %. You may e mail the answers to TLC, info@tlch2o.com or fax the answers to TLC, (928) 272-0747.

This assignment is available to you in a Word Format on TLC's Website. You can find online assistance for this course on the in the Search function on Adobe Acrobat PDF to help find the answers. Student Services (928) 468-0665.

Topic 1 Introduction to Ticks

Multiple choice. Please select one answer only per question. No trick questions.

(S) means the answer may be plural or singular in nature. Or means either answer may work.

1. All ticks have an incomplete metamorphosis: after hatching from the egg a series of similar stages (instars) develop from a_____, to eight legged nymph and then a sexually developed eight legged adult.

A. Six legged larva C. Eight legged larva

B. Seven instar D. None of the above

2. Between each stage there is a molt (ecdysis) which enables the developing tick to expand within a new_____.

A. Idiosoma C. External skeleton

B. Haller's organ D. None of the above

Diet and feeding behaviors

3. Ticks satisfy all of their nutritional requirements as ectoparasites, feeding on a diet of blood in a practice known as _____.

- A. Parasitic mechanisms C. Hematophagy
- B. Hypostome D. None of the above

4. Ticks extract the blood by cutting a hole in the host's epidermis, into which they insert their______, in order to keep the blood from clotting by excreting an anticoagulant or platelet aggregation inhibitor.

- A. Idiosoma C. Hypostome
- B. Coxae D. None of the above

5. While ______, ticks hold onto leaves and grass by their third and fourth pair of legs. They hold the first pair of legs outstretched, waiting to climb on to the host. When a host brushes the spot where a tick is waiting, it quickly climbs onto the host.

A. Questing C. Squatting

B. Aggregation D. None of the above

Genus Ixodes

6. Ticks in the genus Ixodes are easily recognized by the position of the anal groove, which lies in front of the and extends from one side of the body to the other.

A. Capitulum C. Anus

B. Cornua D. None of the above

7. There are 34 species of ticks in the genus Ixodes in the United States, more than in any other genus. Male specimens have a ______ on the ventral side of the abdomen. Most species have enlarged club-like palps.

A. HypostomeB. CornuaC. Complicated arrangement of platesD. None of the above

How a Tick Bites and Feeds

on larval and nymphal ticks are small with less penetration and 8. produce a smaller host reaction.

A. IdiosomaC. MouthpartsB. HypostomeD. None of the above

9. Adult Ixodes and ticks have long mouthparts that can reach the sub dermal layer of skin, produce a larger reaction, and make the tick harder to remove.

- A. Argasidae or Argasid C. Dermacentor
- B. Amblyomma D. None of the above

10. A variety of ______ that aid the feeding process and possibly increase pathogen transmission are introduced in the tick's saliva (e.g., blood platelet aggregation inhibitors, anticoagulants, anti-inflammatory and immunosuppressive enzymes, and vasodilators to increase blood flow).

- A. Vectors of human disease C. Pharmacologically active compounds
- B. Allergic reaction(s)
- D. None of the above

Please complete the entire assignment before submitting the answer key

Topic 2 Tick Identification Section

Tick Life Cvcle

Deer Tick Life Cycle

- 1. The deer tick passes through four life stages (egg, larva, nymph, adult), over a
- A. Two month period C. Two year period
- B. Three month period D. None of the above

Egg to Larvae

2. Their first host is generally a mouse or other medium-sized mammal or bird. Once attached, the larvae embed their mouth parts and feed for several days. If the host is infected with a disease such as Lyme, the tick may be infected during this feeding. The larvae then drop off their host into the leaf litter where they molt into the next stage, the nymph, remaining dormant until the following _____.

- A. Summer C. Spring
- D. None of the above B. Full moon

3	Eggs are fertilized in the fall and deposited in leaf litter the following	They
en	nerge as larvae in late summer of that year, seeking their first blood meal.	

- A. Summer C. Spring
- D. None of the above B. Month

Larvae to Nymph

4. are commonly found on the forest floor in leaf litter and on low lying vegetation. Their host primarily consists of mice and other rodents, deer, birds and unfortunately humans.

- C. Females A. Nymph(s)
- B. Larvae D. None of the above

Nymph to Adult

5.	Over the next few months the nymph molts in	nto the larger ad	lult tick, which e	emerges in fall,
wit	h a peak in October through November.	_	find and feed	on a host, then
the	fomolog lov aggs comptime offer fooding			

the females lay eggs sometime after feeding. A. Both male and female adults C. Larvae

B. Seven instars D. None of the above

Adult Ticks

- 6. Female adults are and larger than males.
- C. Black A. Red or orange
- B. Black and red D. None of the above

7. Adult females infected with disease agents as may transmit disease during this feeding.

- A. Both male and female adults C. Several nymphal stages
- B. Larvae or nymphs D. None of the above
- ticks attach, but do not feed or become engorged. 8.
- A. Nymph(s) C. The adult female
- B. Male D. None of the above

Lone Star Tick Amblyomma americanum

9. Each female produces eggs, which are deposited under leaf and soil litter in middle to late spring.

- A. 300-800
- C. 3,000-8,000
- D. None of the above B. 30,000-80,000

Winter Tick Dermacentor albipictus

is found throughout North America. It is widely distributed throughout 10. California, but populations are concentrated around the central coastal and sierra foothill areas.

- C. This one host tick A. This two host tick
- B. This no host tick D. None of the above

Topic 3 Dangers of Ticks

ticks are removed. A. Allergic Reaction		in humans that is reversible when the
removal of the tick. A. Few weeks	C. Few hours	of the
3. The paralysis may humans when a tick A. Blood	feeds.	transmitted to
parts of the body as A. Disease transmis	f the victim's skull; however	equently associated with the attachment of , the illness occurs from attachment to other ove
Columbia, Canada, a		th America occurs near the border of British ove
and Rocky Mountain A. Lyme disease	spotted fever.	n the United States are
rash, which may dev	elop a characteristic clear c er a tick bite, often in the ar	terized by the development of a large, red entral area (""), ea around the puncture.
 8. The most charact ankles, wrists, and for A. One to two days B. One to two weeks 	C. One to two yea	
	o prevent the transmission ick paralysis is the prompt r C. Hepatitis ses D. None of the ab	emoval of ticks.

10. To remove a tick, grasp it crosswise with narrow tweezers (do not rupture the tick) as close to the point of attachment as possible. _______ tid firmly in the direction of attachment; some back-and-forth wiggling may be necessary. tick

- A. WiggleC. Retract or pullB. RipD. None of the above

Topic 4 Related Blood-feeding Insects

1. Rat control may intensify the attack on humans, but this _____will bite humans even when there is an abundance of host rats on which they can feed. The mite drops from its host after each feeding and may be found on a variety of surfaces near rat-infested areas. It can survive for several days without a blood meal.

- A. Household pest C. Mites
- B. Assassin bugs D. None of the above

House mouse mite (Liponissoides sanguineus)

2. The mite in the U.S. is primarily a parasite of mice. Its major importance is that it has been identified as the vector of ______, a mild and nonfatal human disease.

- A. Grocers' itch
- C. Rat-mite dermatitis
- B. Rickettsial pox D. None of the above

Grain mite

3. Grain mites are reported to have been the cause of mild dermatitis in humans, known under various names as "______," "vanillism" (from infestations on vanilla beans), and "copra itch."

- A. Grocers' itchC. Rickettsial poxB. Rat-mite dermatitisD. None of the above

Life Stages

- 4. Bed bugs may be mistaken for other insects such as , or vice-versa.
- A. Fleas or ticksB. Rodents or RatsC. Booklice and carpet beetlesD. None of the above

Pyrethroids

5. To mimic the insecticidal activity of the natural compound pyrethrum another class of pesticides, pyrethroid pesticides, has been developed. These are_____, which is a sodium channel modulators, and are much less acutely toxic than organophosphates and carbamates. Compounds in this group are often applied against household pests.

- A. Beneficial C. Non-systemic insecticide
- B. Non-persistent D. None of the above

Pyrethroids include:

_____, Bifenthrin, Beta-Cyfluthrin, Cyfluthrin, Cypermethrin, 6. Cyphenothrin, Deltamethrin, Esfenvalerate, Fenpropathrin, Tau-Fluvalinate, Lambda-Cyhalothrin, Gamma Cyhalothrin, Imiprothrin, 1RS cis-Permethrin, Permethrin, Prallethrin, Resmethrin, Sumithrin (d-phenothrin), Tefluthrin, Tetramethrin, Tralomethrin, and Zeta-Cypermethrin

- A. BeneficialC. Non-systemic insecticideB. Allethrin stereoisomersD. None of the above
 - 19

Borates

7. Borax and other sodium bora	tes are used in numerous products such as laundry
additives, eye drops, fertilizers, and	insecticides. Though the mechanisms of toxicity are not
fully understood, boron is	to insects and decay fungi that commonly
damage wood in structures.	
A. An insect growth regulator (IGR)	C. A broad-spectrum pyrethroid insecticide
B. Very toxic	D. None of the above

Common Flea Treatment Pesticides Propoxur

- 8. Propoxur (Baygon®) is a ______insecticide and was introduced in 1959.
- C. Isopropanol A. Carbamate
- B. Sodium channel modulators D. None of the above

The three types of lice that live on humans are:

is known to spread disease. Lice infestations (pediculosis 9. Only the and pthiriasis) are spread most commonly by close person-to-person contact.

- A. Lice C. Body louse
- D. None of the above B. Flea

10. Dogs, cats, and other pets do not play a role in the transmission of .

- A. FleasB. Human liceC. Bed bugsD. None of the above

California DPR Requirement

The Assignment must be submitted to TLC by December 27 in order to be submitted to DPR by the 31st. If it is late, you will be penalized \$50 per day.

When Finished with Your Assignment

REQUIRED DOCUMENTS

Please scan the **Registration Page**, **Proctoring report**, **Answer Key**, Survey and Driver's License and email it to info@TLCH2O.com.

IPhone Scanning Instructions

If you are unable to scan, take a photo of these documents with your **iPhone** and send these photos to TLC, info@TLCH2O.com.

FAX

If you are unable to scan and email, please fax these to TLC, if you fax, call to confirm that we received your paperwork. (928) 468-0675

California DPR Requirement

The Assignment must be submitted to TLC by December 27 in order to be submitted to DPR by the 31st. If it is late, you will be penalized \$50 per day.

Tick Control CEU Training Awareness Assignment #3 For Students Names N-S

You will have 90 days from the start of this course to have successfully passed this assignment with a score of 70 %. You may e mail the answers to TLC, info@tlch2o.com or fax the answers to TLC, (928) 272-0747.

This assignment is available to you in a Word Format on TLC's Website. You can find online assistance for this course on the in the Search function on Adobe Acrobat PDF to help find the answers. Student Services (928) 468-0665.

Topic 1 Introduction to Ticks

Multiple choice. Please select one answer only per question. No trick questions.

(S) means the answer may be plural or singular in nature. Or means either answer may work.

1. After feeding, ticks detach from their host and molt to on the ground, which then feed on larger hosts and molt to adults. Female adults attach to larger hosts, feed, and lay eggs, while males feed very little and occupy larger hosts primarily for mating.

- Ć. The adult female D. None of the above A. Nymph(s)
- B. Larvae

or Soft Ticks contain 193 species, although the composition of the 2. The genera is less certain, and more study is needed before the genera can become stable.

- A. Argasidae or Argasid C. Ornithodoros, and Otobius
- B. Ixodidae D. None of the above

3. feed on blood, and they mate off the host. During feeding, any excess fluid is excreted by the coxal glands, a process which is unique to argasid ticks

A. Nymph(s)

C. The adult female

B. Both male and female adults D. None of the above

Life cycle and reproduction

4. ticks undergo three primary stages of development: larval, nymphal,

and adult.

- A. Only Argasidae or Argasid C. Both ixodid and argasid
- B. Only Dermacentor D. None of the above

Ticks and Mites

5. Ticks of domestic animals directly cause poor health and loss of production to their hosts by many parasitic mechanisms. Ticks also transmit numerous kinds of

between domestic animals. These microbes cause diseases which can be severely debilitating or fatal to domestic animals, and may also affect humans.

- A. Many parasitic mechanisms C. Viruses, bacteria and protozoa
- B. Allergic reaction(s)
- D. None of the above

6. Some mites are parasitic but all ticks are parasitic feeders on blood. Some species of mites may be mistaken for larval ticks at infestations but their _____ are distinctive.

- A. Parasitic mechanismsC. HematophagyB. Feeding mechanismsD. None of the ab D. None of the above

7. All ticks have an incomplete metamorphosis: after hatching from the egg a series of similar stages (instars) develop from a , to eight legged nymph and then a sexually developed eight legged adult.

- A. Six legged larva C. Eight legged larva
- B. Seven instar D. None of the above

8. Ticks find their hosts by detecting animals' breath and body odors, or by sensing body heat, moisture and

C. Vibrations A. Questina B. Aggregation D. None of the above

9. Hard ticks typically take one blood meal in each of the three developmental stages -larval, nymphal and adult. Both sexes are blood feeders, but only the

becomes greatly distended during engorgement. Most species feed on a different host during each stage, but there are some one-host and two-host species.

A. Nymph(s) C. Female

B. Male D. None of the above

10. Male specimens have a ______ on the ventral side of the abdomen. Most species have enlarged club-like palps.

A. HypostomeB. CornuaC. Complicated arrangement of platesD. None of the above

Topic 2 Tick Identification Section

Tick Life Cycle

Deer Tick Life Cycle

- 1. The deer tick passes through four life stages (egg, larva, nymph, adult), over a _____
- A. Two month periodB. Three month periodC. Two year periodD. None of the above

Egg to Larvae

2. Eggs are fertilized in the fall and deposited in leaf litter the following . They emerge as larvae in late summer of that year, seeking their first blood meal. The tiny larva crawls around the forest floor and onto low-lying vegetation looking for an appropriate host.

- A. Summer C. Spring
- B. Month D. None of the above

3. Their first host is generally a mouse or other medium-sized mammal or bird. Once attached, the larvae embed their mouth parts and feed for several days. If the host is infected with a disease such as Lyme, the tick may be infected during this feeding. The larvae then drop off their host into the leaf litter where they molt into the next stage, the nymph,

- remaining dormant until the following
- A. Summer C. Sprina B. Month D. None of the above

Larvae to Nymph

4. ______ are commonly found on the forest floor in leaf litter and on low lying vegetation. Their host primarily consists of mice and other rodents, deer, birds and unfortunately humans.

A. Nymph(s) C. Females

B. Larvae D. None of the above

Nymph to Adult

5. ______find and feed on a host, then the females lay eggs sometime after feeding.

A. Both male and female adults C. Larvae

B. Seven instars D. None of the above

Adult Ticks

- 6. In the fall of the second year, nymphs molt into adult ticks. Female adults are ______and larger than males.
- A. Red or orange C. Black
- B. Black and red D. None of the above

7. Adult females infected with disease agents as _____may transmit disease during this feeding.

- A. Both male and female adults C. Several nymphal stages
- B. Larvae or nymphs D. None of the above

8. _____ticks attach, but do not feed or become engorged. Because the adult males do not take a blood meal, they do not transmit Lyme disease, human anaplasmosis, or babesiosis.

- A. Nymph(s) C. The adult female
- B. Male D. None of the above

Lone Star Tick Amblyomma americanum

9. Each female produces ______ eggs, which are deposited under leaf and soil litter in middle to late spring.

A. 300-800 C. 3,000-8,000

B. 30,000-80,000 D. None of the above

Winter Tick Dermacentor albipictus

10. ______is found throughout North America. It is widely distributed throughout California, but populations are concentrated around the central coastal and sierra foothill areas. It primarily feeds on horses and deer from fall through early spring.

A. This two host tick C. This one host tick

B. This no host tick D. None of the above

Topic 3 Dangers of Ticks

1. Ticks may cause ______in humans that is reversible when the ticks are removed. Symptoms include paralysis of the arms and legs, followed by a general paralysis, which can be fatal if not reversed.

A. Allergic Reaction C. Local infection

B. Paralysis D. None of the above

	cover comp	letely within a		of the
removal of the tick. A. Few weeks	C. Few he	ours		
B. 1 week	D. None o	of the above		
3. The paralysis may humans when a tick fe		by a		transmitted to
A. Blood B. Salivary toxin	D. None of	of the above		
4	the victim'	is fre	quently associated with the illness occurs from a	the attachment of attachment to other
parts of the body as w				
A. Disease transmiss	sion C.			
B. Tick paralysis	D.	None of the abo	Ve	
Columbia, Canada, ar	nd the nort	hwestern	n America occurs near th	
A. United States B. Mexico	C.	Washington		
B. Mexico	D.	None of the abo	Ve	
			the United States are _	
and Rocky Mountain	•			
A. Lyme disease B. Babesiosis				
7. The onset of Lyme	e disease is	s usually characte	erized by the developme entral area ("	
one to two weeks afte	r a tick bite	e, often in the are	a around the puncture.	
A. Bulls eye	C. Cat Sc	ratch Fever	·	
B. Hive	D. None of	of the above		
			ountain spotted fever is a after the	
A. One to two days	C.	One to two year	S	
B. One to two weeks	D.	None of the abo	Ve	
9. The best means to				and
the development of tic A. Bulls Eye Rash		Bis the prompt re Hepatitis	moval of ticks.	
B. Tick-borne disease	es D.	None of the abo	ve	
			row tweezers (do not rup and-forth wiggling may be	
A. Wiggle C. Ret	tract or pull ne of the al			-

Topic 4 Related Blood-feeding Insects

1. Rat control may intensify the attack on humans, but this ______will bite humans even when there is an abundance of host rats on which they can feed. The mite drops from its host after each feeding and may be found on a variety of surfaces near rat-infested areas. It can survive for several days without a blood meal.

A. Household pest C. Mites

B. Assassin bugs D. None of the above

Grain mite

2. Grain mites are reported to have been the cause of mild dermatitis in humans, known under various names as "______," "vanillism" (from infestations on vanilla beans), and "copra itch."

- C. Rickettsial pox A. Grocers' itch
- B. Rat-mite dermatitis D. None of the above

Bed Buas

3. Cimicidae or bed bugs (sometimes bedbugs), are small parasitic insects. The most common type is Cimex lectularius.

A. True B. False

Pyrethroids include:

_, Bifenthrin, Beta-Cyfluthrin, Cyfluthrin, Cypermethrin, 4. Cyphenothrin, Deltamethrin, Esfenvalerate, Fenpropathrin, Tau-Fluvalinate, Lambda-Cyhalothrin, Gamma Cyhalothrin, Imiprothrin, 1RS cis-Permethrin, Permethrin, Prallethrin, Resmethrin, Sumithrin (d-phenothrin), Tefluthrin, Tetramethrin, Tralomethrin, and Zeta-Cvpermethrin

A. Beneficial

- C. Non-systemic insecticide D. None of the above
- B. Allethrin stereoisomers

Permethrin

General Information

5. Permethrin is a ______. It is available in dusts, emulsifiable concentrates, smokes, ULV concentrates, and wettable-powder formulations.

A. An insect growth regulator (IGR) C. A broad-spectrum pyrethroid insecticide

B. Stereoisomers D. None of the above

Common Flea Treatment Pesticides

Propoxur

6. Propoxur (Baygon®) is a ______ insecticide and was introduced in 1959. Propoxur is a non-systemic insecticide with a fast knockdown and long residual effect used against turf, forestry, and household pests and fleas.

A. Carbamate C. Isopropanol

D None of the above B. Sodium channel modulators

Methoprene IGR Treatment

7. Methoprene is _____ with activity against a variety of insect species including horn flies, mosquitoes, beetles, tobacco moths, sciarid flies, fleas (eggs and larvae), fire ants, pharoah ants, midge flies and Indian meal moths.

- A. Not a pyrethrin and permethrin C. An insect growth regulator (IGR)
- B. A stereoisomer D. None of the above

- 8. Dogs, cats, and other pets do not play a role in the transmission of____
- A. Fleas C. Bed bugs
- B. Human lice D. None of the above
- 9. _____ move by crawling; they cannot hop or fly.
- A. Lice C. Assassin bugs
- B. Fleas D. None of the above

Malathion

10. Malathion 0.5% in isopropanol is FDA approved for the treatment of head lice. Apply it to dry hair until the hair and scalp are wet. Leave it on for 12 hours. _____ may be useful for resistant infections.

- A. Malathion C. Non-systemic insecticide
- B. Pyrethrin D. None of the above

California DPR Requirement

The Assignment must be submitted to TLC by December 27 in order to be submitted to DPR by the 31st. If it is late, you will be penalized \$50 per day.

When Finished with Your Assignment

REQUIRED DOCUMENTS

Please scan the **Registration Page**, **Proctoring report**, **Answer Key**, **Survey and Driver's License** and email it to <u>info@TLCH2O.com</u>.

IPhone Scanning Instructions

If you are unable to scan, take a photo of these documents with your **iPhone** and send these photos to TLC, <u>info@TLCH2O.com</u>.

FAX

If you are unable to scan and email, please fax these to TLC, if you fax, call to confirm that we received your paperwork. **(928) 468-0675**

California DPR Requirement

The Assignment must be submitted to TLC by December 27 in order to be submitted to DPR by the 31st. If it is late, you will be penalized \$50 per day.

Tick Control CEU Training Awareness Assignment #4 For Students Names T-Z

You will have 90 days from the start of this course to have successfully passed this assignment with a score of 70 %. You may e mail the answers to TLC, info@tlch2o.com or fax the answers to TLC, (928) 272-0747.

This assignment is available to you in a Word Format on TLC's Website. You can find online assistance for this course on the in the Search function on Adobe Acrobat PDF to help find the answers. Student Services (928) 468-0665.

Topic 1 Introduction to Ticks

Multiple choice. Please select one answer only per question. No trick questions.

(S) means the answer may be plural or singular in nature. Or means either answer may work.

1. More than 800 species of these obligate blood-sucking creatures inhabit the planet. They are second only to mosquitoes as vectors of human disease,

- A. Including parasitic mechanisms
 B. Causing allergic reaction(s)
 C. Both infectious and toxic
 D. None of the above
- B. Causing allergic reaction(s)
- D. None of the above

2. In Ixodidae nymphs and adults, a prominent capitulum (head) projects forwards from the body; in the Argasidae, conversely, the capitulum is concealed beneath the

- A. Idiosoma C. Scutum
- B. Body D. None of the above

3. After feeding, ticks detach from their host and molt to on the ground, which then feed on larger hosts and molt to adults. Female adults attach to larger hosts, feed, and lay eggs, while males feed very little and occupy larger hosts primarily for mating.

- C. The adult female A. Nymph(s)
- B. Larvae D. None of the above

4. Soft ticks have no hard shell (Scutum). In the United States, only ticks of the genus transmit human disease, namely, relapsing fever.

A. Argasidae or Argasid C. Antricola, Argas, Nothaspis, Ornithodoros, and Otobius

B. Ornithodoros D. None of the above

5. feed on blood, and they mate off the host. During feeding, any excess fluid is excreted by the coxal glands, a process which is unique to argasid ticks

C. The adult female A. Nymph(s)

B. Both male and female adults D. None of the above

Ixodidae

- 6. Ixodid ticks require three hosts, and their life cycle takes at least one year to complete. Up to 3,000 eggs are laid on the ground by an adult female tick.
- A. 100 C. 500
- B. 3,000 D. None of the above

7. All ticks have an incomplete metamorphosis: after hatching from the egg a series of similar stages (instars) develop from a _____, to eight legged nymph and then a sexually developed eight legged adult.

- A. Six legged larva C. Eight legged larva
- B. Seven instar D. None of the above

8. Between each stage there is a molt (ecdysis) which enables the developing tick to expand within a new_____.

- A. Idiosoma C. External skeleton
- B. Haller's organ D. None of the above

Argasidae

9. During feeding, any excess fluid is excreted by the_____, a process which is unique to argasid ticks

- A. Idiosoma C. Coxal glands
- B. Haller's organ D. None of the above

General Characteristics and Habits of Hard Ticks (Family Ixodidae)

10. The ______ lie behind the fourth pair of coxae, or basal segments of the leg.

- A. Idiosoma C. Spiracles
- B. Coxae D. None of the above

Please complete the entire assignment before submission of answer key.

Topic 2 Tick Identification Section

Tick Life Cycle

Deer Tick Life Cycle

1. The deer tick passes through four life stages (egg, larva, nymph, adult), over a _____

- A. Two month period C. Two year period
- B. Three month period D. None of the above

Egg to Larvae

2. Eggs are fertilized in the fall and deposited in leaf litter the following_____.

- A. Summer C. Spring
- B. Month D. None of the above

3. The larvae then drop off their host into the leaf litter where they molt into the next stage, the nymph, remaining dormant until the following _____.

- A. Summer C. Spring
- B. Month D. None of the above

Larvae to Nymph

4. During the spring and early summer of the next year the nymphs end their dormancy and begin to seek a host. ______ are commonly found on the forest floor in leaf litter and on low lying vegetation. Their host primarily consists of mice and other rodents, deer, birds and unfortunately humans.

- A. Nymph(s) C. Females
- B. Seven instars D. None of the above

Nymph to Adult

5. Over the next few months the nymph molts into the larger adult tick, which emerges in fall, with a peak in October through Novemberfind and feed on a host, then the females lay eggs sometime after feeding. A. Both male and female adults C. Larvae B. Seven instars D. None of the above
Adult Ticks 6. Female adults areand larger than males. A. Red or orange C. Black B. Black and red D. None of the above
 7. Adult females infected with disease agents asmay transmit disease during this feeding. A. Both male and female adults C. Several nymphal stages B. Larvae or nymphs D. None of the above
 8ticks attach, but do not feed or become engorged. A. Nymph(s) B. Male C. The adult female D. None of the above
Lone Star Tick Amblyomma americanum9. Each female produces eggs, which are deposited under leaf andsoil litter in middle to late spring.A. 300-800C. 3,000-8,000B. 30,000-80,000D. None of the above
Winter Tick Dermacentor albipictus10
Topic 3 Dangers of Ticks 1. Ticks may cause in humans that is reversible when the ticks are removed. A. Allergic Reaction C. Local infection B. Paralysis D. None of the above
 The victim may recover completely within a of the removal of the tick. Few weeks C. Few hours 1 week D. None of the above
 3. The paralysis may be caused by atransmitted to humans when a tick feeds. A. Blood C. Germ B. Salivary toxin D. None of the above

4.

is frequently associated with the attachment of

the tick at the base of the victim's skull; however, the illness occurs from attachment to other parts of the body as well.

A. Disease transmission C. Local infection

D. None of the above B. Tick paralysis

5. The highest incidence of tick paralysis in North America occurs near the border of British Columbia, Canada, and the northwestern

- A. United States C. Washington
- B. Mexico D. None of the above

6. The two most important tick-borne diseases in the United States are and Rocky Mountain spotted fever.

- A. Lyme disease C. Local infections
- B. Babesiosis D. None of the above

7. The onset of Lyme disease is usually characterized by the development of a large, red rash, which may develop a characteristic clear central area ("_____"),

one to two weeks after a tick bite, often in the area around the puncture.

- A. Bulls eye C. Cat Scratch Fever
- B. Hive D. None of the above

8. The most characteristic symptom of Rocky Mountain spotted fever is a rash on the ______after the victim is bitten. C. One to two years ankles, wrists, and forehead

- A. One to two daysC. One to two yearsB. One to two weeksD. None of the above

9. The best means to prevent the transmission of _____ and

the development of tick paralysis is the prompt removal of ticks.

A. Bulls Eye RashC. HepatitisB. Tick-borne diseasesD. None of t

D. None of the above

10. To remove a tick, grasp it crosswise with narrow tweezers (do not rupture the tick) as close to the point of attachment as possible. tick

firmly in the direction of attachment; some back-and-forth wiggling may be necessary.

A. WiggleC. Retract or pullB. Rub itD. None of the above

Topic 4 Related Blood-feeding Insects Mites

1. The straw itch mite and furniture mite come from plant material, and the chigger mite is found in lawns and open woodlands. The tropical rat mite and the mouse mite come from rodents, whereas the itch mite and follicle mite are permanent residents on

A. Household pests C. Humans

B. Rodents or Rats D. None of the above

Tropical rat mite (Ornithonyssus bacoti)

2. This mite is associated with rats throughout the U.S., where it also feeds on humans and many other warm-blooded animals. The bite is painful, causing intense itching and a skin irritation known as _____.

A. Rickettsial pox C. Grocers' itch

B. Rat-mite dermatitis D. None of the above

House mouse mite (Liponissoides sanguineus)

3. The mite in the U.S. is primarily a parasite of mice. Its major importance is that it has been identified as the vector of ______, a mild and nonfatal human disease.

- A. Grocers' itch C. Rat-mite dermatitis
- B. Rickettsial pox D. None of the above

Pyrethroids

4. To mimic the insecticidal activity of the natural compound pyrethrum another class of pesticides, pyrethroid pesticides, has been developed. These are ______, which is a sodium channel modulators, and are much less acutely toxic than organophosphates and carbamates. Compounds in this group are often applied against household pests.

A. Beneficial C. Non-systemic insecticide

B. Non-persistent D. None of the above

Pyrethroids include:

5. ______, Bifenthrin, Beta-Cyfluthrin, Cyfluthrin, Cypermethrin, Cyphenothrin, Deltamethrin, Esfenvalerate, Fenpropathrin, Tau-Fluvalinate, Lambda-Cyhalothrin, Gamma Cyhalothrin, Imiprothrin, 1RS cis-Permethrin, Permethrin, Prallethrin, Resmethrin, Sumithrin (d-phenothrin), Tefluthrin, Tetramethrin, Tralomethrin, and Zeta-Cypermethrin

A. Beneficial

C. Non-systemic insecticide

B. Allethrin stereoisomers D. None of the above

Permethrin

General Information

6. Permethrin is a ______. It is available in dusts, emulsifiable concentrates, smokes, ULV concentrates, and wettable-powder formulations.

A. An insect growth regulator (IGR) C. A broad-spectrum pyrethroid insecticide

B. Stereoisomers D. None of the above

Borates

7. Borax and other sodium borates are used in numerous products such as laundry additives, eye drops, fertilizers, and insecticides. Though the mechanisms of toxicity are not fully understood, boron is _______to insects and decay fungi that commonly damage wood in structures.

A. An insect growth regulator (IGR)B. Very toxicC. A broad-spectrum pyrethroid insecticideD. None of the above

Common Flea Treatment Pesticides

Propoxur

- 8. Propoxur (Baygon®) is a ______ insecticide and was introduced in 1959.
- A. Carbamate C. Isopropanol
- B. Sodium channel modulators D. None of the above

Malathion

9. Malathion 0.5% in isopropanol is FDA approved for the treatment of head lice. Apply it to dry hair until the hair and scalp are wet. Leave it on for 12 hours. _____ may be useful for resistant infections.

A. Malathion C. Non-systemic insecticide

B. Pyrethrin D. None of the above

Lindane

10. Lindane has been associated with ______suffered both by people being treated and by people applying the treatment. It is also a troublesome pollutant of wastewater and requires special treatment to be removed.

A. A variety of adverse reactions C. An insect growth regulators

B. Long residual effects D. None of the above

California DPR Requirement

The Assignment must be submitted to TLC by December 27 in order to be submitted to DPR by the 31st. If it is late, you will be penalized \$50 per day.

When Finished with Your Assignment

REQUIRED DOCUMENTS

Please scan the **Registration Page, Proctoring report, Answer Key, Survey and Driver's License** and email it to <u>info@TLCH2O.com</u>.

IPhone Scanning Instructions

If you are unable to scan, take a photo of these documents with your **iPhone** and send these photos to TLC, <u>info@TLCH2O.com</u>.

FAX

If you are unable to scan and email, please fax these to TLC, if you fax, call to confirm that we received your paperwork. **(928) 468-0675**

California DPR Requirement

The Assignment must be submitted to TLC by December 27 in order to be submitted to DPR by the 31st. If it is late, you will be penalized \$50 per day.

Tick Control CEU Training Awareness Assignment #5 Alternative Assignment for Repeat Students

You will have 90 days from the start of this course to have successfully passed this assignment with a score of 70 %. You may e mail the answers to TLC, info@tlch2o.com or fax the answers to TLC, (928) 272-0747.

This assignment is available to you in a Word Format on TLC's Website. You can find online assistance for this course on the in the Search function on Adobe Acrobat PDF to help find the answers. Student Services (928) 468-0665.

Topic 1 Introduction to Ticks

Multiple choice. Please select one answer only per question. No trick questions.

(S) means the answer may be plural or singular in nature. Or means either answer may work.

1. Both sexes are blood feeders, but only the ______ becomes greatly distended during engorgement. Most species feed on a different host during each stage, but there are some one-host and two-host species.

- A. Nymph(s) C. Female
- B. Male D. None of the above

2. The lone star tick, Amblyomma americanum, can carry spotted fever, tularemia, and possibly Q fever. Female specimens are easily recognized by the conspicuous silvery-white spot at the tip of the______, hence the name "speck-back" in the Ozark Mountains and the common name "lone star tick" for the Lone Star State of Texas.

- A. Idiosoma C. Scutum
- B. Capitulum D. None of the above

3. More than 800 species of these obligate blood-sucking creatures inhabit the planet. They are second only to mosquitoes as vectors of human disease, _____.

- A. Including parasitic mechanisms C. Both infectious and toxic
- B. Causing allergic reaction(s)
 - eaction(s) D. None of the above

4. Ixodid ticks require three hosts, and their life cycle takes at least one year to complete. Up to 3,000 eggs are laid on the ground by an adult female tick.

- A. 100 C. 500
- B. 3,000 D. None of the above

5. Ticks, like mites, have bodies which are divided into two primary sections: the anterior capitulum (or gnathosoma), which contains the head and mouthparts; and the posterior which contains the legs, digestive tract, and reproductive organs.

- A. Ediosoma C. Idiosoma
- B. Hypostome D. None of the above

6. The Gulf Coast tick, Amblyomma maculatum, is found particularly along the Gulf and South Atlantic coastlines. It has spurs on the _____ and more diffuse pale markings on the female than does the lone star tick.

A. True B. False

7. Ticks satisfy all of their nutritional requirements as ectoparasites, feeding on a diet of blood in a practice known as _____. They are obligate hematophages, needing blood to survive and move from one stage of life to another.

- A. Parasitic mechanisms C. Hematophagy
- D. None of the above B. Hypostome

8. While _____, ticks hold onto leaves and grass by their third and fourth pair of legs. They hold the first pair of legs outstretched, waiting to climb on to the host. When a host brushes the spot where a tick is waiting, it guickly climbs onto the host.

- A. QuestingC. SquattingB. AggregationD. None of the above
- ticks undergo three primary stages of development: larval, nymphal, 9.
- and adult.
- A. Only Argasidae or ArgasidB. Only DermacentorC. Both ixodid and argasidD. None of the above

10. In the female, the ______ covers only a part of the dorsal surface and is almost obscured when she becomes engorged.

- A. Capitulum C. Scutum
- B. Coxae D. None of the above

Topic 2 Tick Identification Section

- 1. The deer tick passes through four life stages (egg, larva, nymph, adult), over a
- A. Two month periodČ. Two year periodB. Three month periodD. None of the above
- 2. Eggs are fertilized in the fall and deposited in leaf litter the following .
- A. Summer C. Spring
- D. None of the above B. Month

3. The larvae then drop off their host into the leaf litter where they molt into the next stage, the nymph, remaining dormant until the following

- A. Summer C. Spring
- B. Month D. None of the above

4. During the spring and early summer of the next year the nymphs end their dormancy and begin to seek a host. ______ are commonly found on the forest floor in leaf litter and on low lying vegetation.

- A. Nymph(s)
- C. Females
- A. Nymph(s)C. FemalesB. Seven instarsD. None of the above

5. find and feed on a host, then the females lay eggs sometime after

feeding.

- A. Both male and female adults C. Larvae
- B. Seven instars D. None of the above
- 6. In the fall of the second year, nymphs molt into adult ticks. Female adults are and larger than males.
- A. Red or orange C. Black
- B. Black and red D. None of the above

7. As female ticks feed over the course of several days, their bodies slowly enlarge with blood (engorge). Adult females infected with disease agents as may transmit disease during this feeding.

A. Both male and female adults C. Several nymphal stages

B. Larvae or nymphs D. None of the above

ticks attach, but do not feed or become engorged. Because the adult 8. males do not take a blood meal, they do not transmit Lyme disease, human anaplasmosis, or babesiosis.

- A. Nymph(s) C. The adult female
- B. Male D. None of the above

9. Each female produces ______ eggs, which are deposited under leaf and soil litter in middle to late spring.

A. 300-800 C. 3,000-8,000

B. 30,000-80,000 D. None of the above

is found throughout North America. It is widely distributed throughout 10. California, but populations are concentrated around the central coastal and sierra foothill areas. It primarily feeds on horses and deer from fall through early spring.

C. This one host tick A. This two host tick

B. This no host tick D. None of the above

Topic 3 Dangers of Ticks

in humans that is reversible when the 1. Ticks may cause ticks are removed. Symptoms include paralysis of the arms and legs, followed by a general paralysis, which can be fatal if not reversed.

- A. Allergic Reaction C. Local infection
- B. Paralvsis D. None of the above

2. The victim may recover completely within a of the removal of the tick.

- A. Few weeks C. Few hours
- D. None of the above B. 1 week
- 3. The paralysis may be caused by a ______transmitted to humans when a tick feeds.
- A. Blood C. Germ
- B. Salivary toxin D. None of the above

4.

is frequently associated with the attachment of

the tick at the base of the victim's skull; however, the illness occurs from attachment to other parts of the body as well.

- A. Disease transmission C. Local infection
- B. Tick paralysis D. None of the above

5. The highest incidence of tick paralysis in North America occurs near the border of British Columbia, Canada, and the northwestern_____.

- A. United States C. Washington
- B. Mexico D. None of the above

6. The two most important tick-borne diseases in the United States are ______ and Rocky Mountain spotted fever.

- A. Lyme disease C. Local infections
- B. Babesiosis D. None of the above

7. The onset of Lyme disease is usually characterized by the development of a large, red rash, which may develop a characteristic clear central area ("_____"),

one to two weeks after a tick bite, often in the area around the puncture.

- A. Ring o C. Local infection
- B. Bulls eye D. None of the above

8. The most characteristic symptom of Rocky Mountain spotted fever is a rash on the ankles, wrists, and forehead after the victim is bitten.

- A. One to two days C. One to two years
- B. One to two weeks D. None of the above
- 9. The best means to prevent the transmission of ______and

the development of tick paralysis is the prompt removal of ticks.

- A. Bulls Eye Rash C. Hepatitis
- B. Tick-borne diseases D. None of the above

10. To remove a tick, grasp it crosswise with narrow tweezers (do not rupture the tick) as close to the point of attachment as possible. ________ tick firmly in the direction of attachment; some back-and-forth wiggling may be necessary.

A. Wiggle C. Retract or pull

B. Break Off D. None of the above

Topic 4 Related Blood-feeding Insects

1. To mimic the insecticidal activity of the natural compound pyrethrum another class of pesticides, pyrethroid pesticides, has been developed. These are ______, which is a sodium channel modulators, and are much less acutely toxic than organophosphates and carbamates. Compounds in this group are often applied against household pests.

A. Beneficial C. Non-systemic insecticide

B. Non-persistent D. None of the above

2. The tropical rat mite and the mouse mite come from rodents, whereas the itch mite and follicle mite are permanent residents on _____.

- A. Household pests C. Humans
- B. Rodents or Rats D. None of the above

3
B. Flying ticks D. None of the above
 4 do not have hind legs to hop or jump. Head lice do not have wings and cannot fly. A. Head lice C. Fleas B. Flying ticks D. None of the above
 5. This mite is associated with rats throughout the U.S., where it also feeds on humans and many other warm-blooded animals. The bite is painful, causing intense itching and a skin irritation known as A. Rickettsial pox C. Grocers' itch B. Rat-mite dermatitis D. None of the above
 6. The mite in the U.S. is primarily a parasite of mice. Its major importance is that it has been identified as the vector of, a mild and nonfatal human disease. A. Grocers' itch B. Rickettsial pox C. Rat-mite dermatitis D. None of the above
 7. Cimicidae or bed bugs are small parasitic insects. The most common type is Cimex lectularius. A. True B. False
 8. Bed bugs may be mistaken for other insects such as, or vice-versa. A. Fleas or ticks C. Booklice and carpet beetles B. Assassin bugs D. None of the above
 9. Permethrin is a It is available in dusts, emulsifiable concentrates, smokes, ULV concentrates, and wettable-powder formulations. A. An insect growth regulator (IGR) C. A broad-spectrum pyrethroid insecticide B. Stereoisomers D. None of the above
Common Flea Treatment Pesticides Propoxur 10. Propoxur (Baygon®) is a insecticide and was introduced in 1959. Propoxur is a non-systemic insecticide with a fast knockdown and long residual effect used against turf, forestry, and household pests and fleas. It is also used in pest control for other domestic animals. A. Carbamate C. Isopropanol B. Sodium channel modulators D. None of the above

California DPR Requirement

The Assignment must be submitted to TLC by December 27 in order to be submitted to DPR by the 31st. If it is late, you will be penalized \$50 per day.